



Correction: Folic acid conjugated with serum albumin for nano- and submicron delivery systems for applications in therapy and diagnostics

Maria G. Gorobets^{1*} , Anna V. Toroptseva¹ , Madina I. Abdullina¹ , Vadim S. Pokrovsky^{1,2,3} ,
Derenik S. Khachatryan¹ , Anna V. Bychkova¹ 

¹Emanuel Institute of Biochemical Physics of Russian Academy of Sciences, 119334 Moscow, Russia

²N.N. Blokhin National Medical Research Center of Oncology, 115478 Moscow, Russia

³Laboratory of Experimental Oncology, Research Institute of Molecular and Cellular Medicine, RUDN University, 117198 Moscow, Russia

***Correspondence:** Maria G. Gorobets, Emanuel Institute of Biochemical Physics of Russian Academy of Sciences, 4, Kosygina Str., 119334 Moscow, Russia. maria.g.gorobets@gmail.com

Published: September 24, 2025

Cite this article: Gorobets MG, Toroptseva AV, Abdullina MI, Pokrovsky VS, Khachatryan DS, Bychkova AV. Correction: Folic acid conjugated with serum albumin for nano- and submicron delivery systems for applications in therapy and diagnostics. *Explor Drug Sci.* 2025;3:1008130. <https://doi.org/10.37349/eds.2025.1008130>

Correction to “Folic acid conjugated with serum albumin for nano- and submicron delivery systems for applications in therapy and diagnostics” by Gorobets MG, Toroptseva AV, Abdullina MI, Pokrovsky VS, Khachatryan DS, Bychkova AV. *Explor Drug Sci.* 2025;3:1008101. DOI: [10.37349/eds.2025.1008101](https://doi.org/10.37349/eds.2025.1008101)

In the recently published article, an error was identified in the description of the molecular interactions (in the section “Conjugation of FA residue to serum albumin”). The article incorrectly stated that “For a mixture of bovine albumin and NHS-ester of FA, it was shown an engagement in interaction via different mechanisms of the following listed in the brackets amino acid residues: hydrogen bonding (Tyr149, Tyr340, Glu152, Glu339, Gln220, His289, Val342), π - π interaction (Tyr156), hydrophobic interaction (Lys187, Arg194, Arg256).” The correct statement is that “For a mixture of bovine albumin and NHS-ester of FA, it was shown an engagement in interaction via different mechanisms of the following listed in the brackets amino acid residues: hydrogen bonding (Tyr149, Tyr340, Glu152, Glu339, Gln220, His289, Val342), π - π and/or hydrophobic interactions (Tyr156, Lys187, Arg194, Arg256).” This has now been amended in the HTML and PDF versions of the article.

The original uncorrected PDF can be accessed from <https://www.explorationpub.com/uploads/er1008101.pdf>

