







Application of fuzzy logic techniques for sensory evaluation of plant-based extrudates fortified with bioactive compounds

Mekala Pavani¹, Poonam Singha¹, Darwin Thanaraj Rajamanickam², Sushil Kumar Singh^{1*}

¹Department of Food Process Engineering, National Institute of Technology, Odisha 769008, India

²Vetio Animal Health, Jupiter, FL 33478, USA

***Correspondence:** Sushil Kumar Singh, Department of Food Process Engineering, National Institute of Technology, Sector 1, Rourkela, Odisha 769008, India. sksingh32325@gmail.com; singhsk@nitrrkl.ac.in

Academic Editor: Di Wu, Queen's University of Belfast, UK

Received: July 21, 2023 **Accepted:** November 23, 2023 **Published:** December 29, 2023

Cite this article: Pavani M, Singha P, Rajamanickam DT, Singh SK. Application of fuzzy logic techniques for sensory evaluation of plant-based extrudates fortified with bioactive compounds. *Explor Foods Foodomics*. 2023;1:272–87. <https://doi.org/10.37349/eff.2023.00021>

Abstract

Aim: This study aims to evaluate the sensory profile of corn-based extrudates fortified with phytosterol and pea protein isolates (PPI) using the fuzzy logic technique to assess similarity values and rank the quality attributes.

Methods: Using a mix of yellow PPI (ranging from 0 to 20%) and corn flour (ranging from 80% to 100%), extrudates were developed, ensuring a consistent addition of phytosterol-containing oil at 5%. For this experiment, the Box-Behnken (BB) design was used, comprising 17 runs, factoring in parameters like the percentage of PPI (0–20%), screw speed (300–500 rpm), and temperature (130°–150°C). The optimal conditions were found to be 2.78% PPI, a screw speed of 451 rpm, and a temperature of 150°C, resulting in a desirability value of 0.725. For sensory evaluation, the fuzzy logic technique was used to compare the functional extrudates (S_1) with commercial variants (S_2 , S_3 , and S_4). This helped to gauge acceptance/rejection, similarity values, rankings, and overall consumer acceptability of the extrudates.

Results: Commercial sample S_4 achieved the highest ranking on the sensory scale as “very good”. When considering the quality attributes of extrudates, taste and mouthfeel were the most favored, followed by color and flavor. This study underscored the value of using fuzzy logic for sensory evaluation in determining the acceptance of new food products. It also proved effective in assessing food products' quality attributes, especially after evaluating the phytosterol content post-extrusion.

Conclusions: The fuzzy logic technique in sensory evaluation has effectively identified the optimal extrudates and their quality attributes during the development of new functional food.

Keywords

Fuzzy logic, functional extrudates, phytosterols, pea protein isolate



Introduction

In recent years, the demand for ready-to-eat snacks has risen, reflecting shifts in human dietary habits and lifestyles. As consumers grow more health-conscious, the methods used to create snack products have evolved. Among these methods, extrusion cooking technology stands out for its cost-effectiveness, ability to produce snacks in diverse shapes and sizes, reduced energy consumption, and convenience [1–3]. It further contributes to starch gelatinization and protein denaturation [4, 5]. Many snacks in the market are from starch-based sources such as corn, wheat, rice, potato, cassava, and sweet potato [6–9]. However, these starch-based sources are nutritionally poor in terms of protein, minerals, fibers, and bioactive compounds. Additionally, overconsumption of fried starch-based snacks can lead to health issues such as obesity, high blood pressure, cancer, and elevated glycemic levels [10]. Hence, both food scientists and industry are exploring alternatives. This includes combining existing starch-based snacks with plant-based proteins [11, 12], fortification with bioactive compounds, and by-products [13, 14] to enhance nutritional value and mitigate health issues.

Bioactive compounds mainly find application in the development of newer food products, medicine, agriculture, and industry [15]. Phytosterols are plant-based sterol fractions gaining importance in various food applications due to their cholesterol-lowering properties, which aids in lowering hypercholesterolemia in human blood serum [16, 17]. They are naturally extracted from oil seeds, fruits, vegetables, cereals, nuts, and microalgae, but they are available in much less quantities, i.e., 180–400 mg. Several studies have reported that consumption of 3 g of phytosterol per day could help to lower the cholesterol level by 12% [18]. Therefore, they must be supplied through dietary supplements, pills, and fortified foods to provide numerous health benefits [19]. However, the application of phytosterols in various food products is still challenging due to their lower solubility (in water and oil), crystalline behavior, and lower bioavailability. Till now, various dairy products, fruit juices, etc., have been fortified with phytosterols. However, studies on fortification of extrudates with phytosterols are scarce [20].

This study will investigate the impact of adding bioactive compounds (phytosterols) and plant-based proteins [pea protein isolates (PPI)] to corn-based extrudates. Along with phytosterol, PPI is incorporated due to its better nutritional properties, greater sustainability, lower cost, non-allergic, and gluten-free nature [21, 22]. It contains around 20–30% protein based on the extraction procedure, genotype, and variety. The use of PPI in various food applications has been gaining importance. For example, Beck et al. [23], Philipp et al. [24], and Philipp et al. [25] studied the fortification of rice-based extrudates with PPI.

During the extrusion process, the raw materials containing carbohydrate, protein, and fat undergo a combination of thermo-mechanical stresses resulting in changes in physicochemical, textural, structural properties, and sensory characteristics of the developed extrudates [6, 7, 26–30]. Most of the properties except sensory evaluation can be measured using standard procedure using different instruments or standard methods. Sensory appeal, particularly taste is an ultimate criterion for consumers for acceptance or rejection of novel food products. Traditional techniques have been used for sensory analysis that can only be perceived by sight, smell, in terms of qualitatively but are unable to provide the analysis in the precise way. Experimental procedures are often too expensive, time-consuming, and may lack a generalized theoretical description of the process [31, 32]. Fuzzy logic technique could be an option, in which arbitrary and subjective data can be analyzed to derive important conclusions regarding acceptance/rejection, strong/weak attributes, and ranking of food products by using single or multiple panelists [33]. Till now, numerous studies have been conducted on the sensory evaluation of food products using fuzzy logic techniques. These include chana poda [34], soup mix [35], millet-based bread enriched with dietary fiber [36], beverages [37], mango beverages [38], litchi drinks [39], milk containing barberry juice [40] and many other food products. In this study, sensory assessment of functional extrudates using the fuzzy logic approach has been reported.

Materials and methods

Raw materials

The corn seeds, soybean oil, commercial extruded snacks, and seasonings were procured from the local market in Rourkela, Odisha, India. Yellow PPI was purchased from MyFitFuel Pvt. Ltd., India, and phytosterol (predominantly containing 75% beta-sitosterol and 10% campesterol) was purchased from Thermo Scientific Pvt. Ltd., India.

Dispersion of phytosterols

In this study, soybean oil was selected for phytosterol dispersion based on the previous study conducted by Pavani et al. [41]. Initially, oil was heated at 90°C, and phytosterol in the solid form (1%, w/v) was added into the oil and continuously stirred for 15 mins until phytosterols were completely dissolved in the oil. The resulting oil was considered a functional oil throughout the study.

Development of extrudates containing phytosterols

Prior to extrusion, the composite blends (I, II, and III) were prepared as shown in Table 1. The blends were mixed and maintained at a moisture content of 18%, wet basis. The hydrated blends were kept overnight at room temperature (~25°C) in air-tight containers before the extrusion experiments. A laboratory-scale twin screw extruder (Jinan Saibainuo Machinery Co., Ltd.) with a circular orifice die, and an aperture diameter of 3 mm was used for the extrusion process. The operating power was 14.5 kW, and the barrel length (L) to diameter (D) ratio of the extruder was 24.5. The extruder barrel consists of four zones as shown in Figure 1. The first three zones' temperatures (T_1 , T_2 , T_3 from left to right side as shown in Figure 1) were maintained at temperatures of 30°C, 50°C, and 70°C, respectively, whereas the fourth zone temperature T_4 was varied from 130°C to 150°C. BB design was used with PPI, screw speed, and temperature of zone four (T) as process parameters (Table 2). A total of 17 experimental runs were carried out with five center points. The extrudates collected after each experiment were dried using a hot air oven at 40°C for 3 h or until constant weight was reached. The extrudates collected at optimum conditions (PPI of 2.78%, screw speed of 451 rpm, and T of 150°C) with a desirability value of 0.725 were selected for conducting sensory studies and are referred to as the functional extrudate (S_1) [42].

Phytosterol retention in functional extrudates

The phytosterol retention (PR) of the functional extrudates was calculated using the procedure of Araújo et al. [43]. Before analysis, the hydroethanolic extract was prepared by maceration of extruded powder with a solvent ratio of 2:10 (%; w/v) for 7 days. Liebermann-Burchard reagent was prepared by mixing 5 mL of sulfuric acid into acetic anhydride (50 mL). 100 mL of hydroethanolic extract which was initially dried at 40°C. Then, the residue was resuspended into 20 mL of chloroform, and the volume was adjusted to 50 mL using the same solvent. From the above solution, 10 mL was added with 2 mL of LB reagent, and absorbance was measured at 625 nm after adding the reagent. The phytosterol content was measured using equation 1:

$$PR\left(\frac{g}{100g}\right) = Cs \times \frac{A_a}{A_s}$$

C_s is the standard concentration, A_a is the absorbance of the sample, and A_s is the absorbance of the standard.

Training of the panelists/judges

Before sensory evaluation, all panel members were requested to complete training that was provided on various sensory evaluation terminologies, sampling techniques, sensory scale scoring, selection of quality attributes, sensory scale, different scoring methods, and data interpretation to turn each response into a score.

Table 1. Ingredient composition of feed blends

Feed composition	Mass of the ingredients (g kg ⁻¹)		
	Blend I	Blend II	Blend III
PPI	0	100	200
Corn flour	950	850	750
Functional oil	50	50	50




Figure 1. Schematic diagram of the twin screw extruder [42]

Note. Adapted with permission from “Impact of extrusion processing on bioactive compound enriched plant-based extrudates: a comprehensive study and optimization using RSM and ANN-GA,” by Pavani M, Singha P, Rajamanickam DT, Singh SK. *Future Foods*. 2023;100286 (<https://doi.org/10.1016/j.fufo.2023.100286>). © 2023 The Author(s).

Table 2. Independent variables and their levels

Independent variables	Coded values		
	−1	0	+1
PPI (%)	0	10	20
Screw speed (rpm)	300	400	500
Temperature (°C)	130	140	150

Sensory evaluation

Sensory evaluation of functional extrudates (S_1) and commercial samples (S_2 , S_3 , and S_4) were conducted to understand the acceptance or rejection of extrudates, similarity values, ranking, and consumer acceptability of the extrudates. Henceforth, functional extrudates (S_1) and commercial samples (S_2 , S_3 , and S_4) will be referred to as “samples” wherever they are mentioned together. The evaluation was conducted with fifteen panelists (nine men and six women between the ages of 25 and 35) from NIT Rourkela, Odisha, India. They were selected based on their interest in sensory evaluation, familiarity with snack products, and willingness to participate in the sensory evaluation. Panelists reported having no illness during the sensory evaluation. Before the experiment, extrudates were placed on paper plates and named randomly. Each panelist was asked to taste the extrudates and provide their preference by a tick mark (✓) on the linguistic score sheet. Panelists rinsed their mouth with water between assessments of various extrudates to prevent fatigue and lingering taste.

Fuzzy logic analysis

The relationship between independent attributes, i.e., quality characteristics (color, flavor, taste, and mouthfeel), and output variables, i.e., panelist’s preference (acceptance or rejection, similarity values, and ranking), is numerically measured using the fuzzy logic technique. In this process, the score obtained from each panelist was mathematically converted and interpreted into a triangular membership function (MF). Then, each sample and its corresponding quality features were ranked in general and separately. The flow chart (Figure 2) shows the various steps involved in the sensory evaluation of extrudates using the fuzzy

logic technique. Triplets of each sample and their corresponding quality attributes were calculated using a sensory scale followed by the overall sensory scores (SS) of samples and quality attributes in general. Next, overall MF values were calculated using a standard fuzzy scale (F). Finally, the similarity values, the ranking of the samples, and their quality parameters were measured.




Figure 2. Process flow for various steps involved in the sensory evaluation of the samples

Triplets associated with sensory scales

Triangular MF distribution pattern of sensory scales is represented by a set of three numbers, called “triplet” [44]. Values of triplets associated with triangular membership distribution function for five-point sensory scale are shown in Figure 3. Δabc represents membership distribution function for the not satisfactory/not at all important (NI) category, Δac_1d represents distribution function for fair/somewhat important (SI) category, etc. The first number of the triplet denotes the coordinate of abscissa at which the value of MF is 1. The second and third numbers of the triplet designate the distance to left and right respectively of the first number where the MF is zero.

Triplets for SS of the samples and quality attributes

The triplets for SS of the individual sample were assessed in terms of: (i) the sum of SS; (ii) triplets associated with the sensory scale; and (iii) number of panelists as shown in equation 2:

$$S(r, \alpha) = \frac{n_1(0 \ 0 \ 25) + n_2(25 \ 25 \ 25) + n_3(50 \ 25 \ 25) + n_4(75 \ 25 \ 25) + n_5(100 \ 25 \ 0)}{n_1 + n_2 + n_3 + n_4 + n_5}$$

Where S indicates the triplets for the SS for particular samples (r) for a particular quality attribute (α); r denotes the samples (S_1, S_2, S_3, S_4); α denotes sample’s quality attributes, i.e., color (C), flavor (A), taste (T), and mouthfeel (M); n_1, n_2, n_3, n_4 , and n_5 are the number of judges who rated particular sample (r) as unsatisfactory, fair, medium, good, and excellent, respectively on a five-point sensory scale.




Figure 3. Values of triplets associated with triangular membership distribution function for 5-point sensory scale [44]

Note. Adapted with permission from "Sensory evaluation using fuzzy logic," by Das H. In: Food processing operations analysis. New Delhi: Asian Books Pvt. Ltd.; 2005. pp. 383–402. © 2005 Publisher.

Similarly, for a particular quality attribute (color, flavor, taste, and mouthfeel) of the samples in general, the values of triplets QC , QA , QT , and QM were calculated using equation 3:

$$Q\beta = \frac{n_1(0 \ 0 \ 25) + n_2(25 \ 25 \ 25) + n_3(50 \ 25 \ 25) + n_4(75 \ 25 \ 25) + n_5(100 \ 25 \ 0)}{n_1 + n_2 + n_3 + n_4 + n_5}$$

Where β denotes quality attribute, i.e., color (C), flavor (A), taste (T), and mouthfeel (M); n_1 , n_2 , n_3 , n_4 , and n_5 represents the number of judges opting for specific linguistic scores for the quality attributes in general.

Triplets for relative weights of the quality attributes

The triplets for the relative weights of the quality attributes were determined to find out the triplets for the overall SS of the samples. Equation 4 was used to calculate triplets for the relative weights for color:

$$QC_{rel} = \frac{QC}{Q_{sum}}$$

Where Q_{sum} is the sum of the first digits of triplets of all the quality attributes, i.e., QC , QA , QT , and QM , whereas QC_{rel} is triplet for the relative weights for the color. Similarly, triplets for the relative weights for the flavor (QA_{rel}), taste (QT_{rel}), and mouthfeel (QM_{rel}) were calculated.

Triplets for overall scores of the samples

Overall SS of S_1 were calculated using equation 5:

$$SO_1 = (S_1C \times QC_{rel}) + (S_1A \times QA_{rel}) + (S_1T \times QT_{rel}) + (S_1M \times QM_{rel})$$

Where SO_1 specifies the overall SS of S_1 ; S_1C , S_1A , S_1T , and S_1M are triplets of the SS of S_1 related to color, flavor, taste, and mouthfeel, respectively; QC_{rel} , QA_{rel} , QT_{rel} , and QM_{rel} indicates the triplet for the relative weight of the quality attributes of the samples in general. Each term on the right-hand side of equation 5 is a triplet, and the multiplication of the two triplets for e.g., triplet ($p \ q \ r$) and triplet ($s \ t \ u$) is done by following equation 6:

$$(p \ q \ r) \times (s \ t \ u) = (p \times s \ p \times t + s \times q \ p \times u + s \times r)$$

Where triplet is a combination of three integers which is used to represent the distribution patterns of triangular MF. Using the similar procedure, overall SS for other samples, i.e., SO_2 , SO_3 , and SO_4 were determined.

Computation of overall MF function of SS on the standard fuzzy scale

The triangular distribution pattern of 6-point sensory scale, which will be referred here as “standard fuzzy scale” is shown in Figure 4. The standard fuzzy scale represents six sensory scales namely F_1 , F_2 , F_3 , F_4 , F_5 , and F_6 where the linguistic terminology for the scale factors has been modified to fit the case of sensory analyses. MF of each of the sensory scales follows triangular distribution pattern where maximum value of membership is 1. Values of MF of F_1 through F_6 are defined by a set of 10 numbers, which are defined in equation 7:

$$\begin{aligned} F_1 &= (1, 0.5, 0, 0, 0, 0, 0, 0, 0, 0) \\ F_2 &= (0.5, 1, 1, 0.5, 0, 0, 0, 0, 0, 0) \\ F_3 &= (0, 0, 0.5, 1, 1, 0.5, 0, 0, 0, 0) \\ F_4 &= (0, 0, 0, 0, 0.5, 1, 1, 0.5, 0, 0) \\ F_5 &= (0, 0, 0, 0, 0, 0, 0.5, 1, 1, 0.5) \\ F_6 &= (0, 0, 0, 0, 0, 0, 0, 0, 0.5, 1) \end{aligned}$$




Figure 4. Triangular membership distribution for 6-point sensory scale “standard fuzzy scale” [44]. Values of F_1 to F_6 are defined by a set of 10 numbers, which are the maximum numbers between each two consecutive points from 0 to 100, which could be defined as “(maximum value of MF observed between 0 and 10), (maximum value of MF observed between 10 and 20), (maximum value of MF observed between 20 and 30), (maximum value of MF observed between 30 and 40), (maximum value of MF observed between 40 and 50), (maximum value of MF observed between 50 and 60), (maximum value of MF observed between 60 and 70), (maximum value of MF observed between 70 and 80), (maximum value of MF observed between 80 and 90), and (maximum value of MF observed between 90 and 100)”

Note. Adapted with permission from “Sensory evaluation using fuzzy logic,” by Das H. In: Food processing operations analysis. New Delhi: Asian Books Pvt. Ltd.; 2005. pp. 383–402. © 2005 Publisher.

Now the values of the MF of the overall SS of samples were determined on the standard fuzzy scale using the triplets of the SS calculated from equation 5. The triplet is a combination of three integers (a b c), which is used to represent the distribution patterns of triangular MF across sensory scale as shown in Figure 5. The value of MF is 1 when the abscissa x is “ a ” and zero when x is less than ($a - b$) or greater than ($a + c$).

For a given value of x on abscissa, value of MF, B_x was calculated using equation 8:

$$\begin{aligned} B_x &= \frac{x - (a - b)}{b} \quad \text{for } (a - b) < x < a \\ &= \frac{(a + c) - x}{c} \quad \text{for } a < x < (a + c) \\ &= 0 \quad \text{for all other values of } x \end{aligned}$$

$$\begin{aligned} B_x &= 0 \quad \text{for } x < (a - b) \quad \& \quad x > (a + c) \\ B_x &= 1 \quad \text{at } x = a \end{aligned}$$




Figure 5. Graphical representation of the triplets ($a\ b\ c$) and its MF [44]

Note. Adapted with permission from “Sensory evaluation using fuzzy logic,” by Das H. In: Food processing operations analysis. New Delhi: Asian Books Pvt. Ltd.; 2005. pp. 383–402. © 2005 Publisher.

For each of the samples and its triplet associated with overall SS, the value of MF B_x at $x = 0, 10, 20, 30, 40, 50, 60, 70, 80, 90$, and 100 was calculated using equation 8. The value of the MF of the overall SS of each sample on the standard fuzzy scale is also determined by a set of 10 numbers, which are the maximum values of B_x in the 10 intervals from 0 to 100 in the mentioned range of x between $(a - b)$ and $(a + c)$. Hence, ten intervals, where membership values of B_x are calculated, comprise “ $0 < x < 10, 10 < x < 20, 20 < x < 30, 30 < x < 40, 40 < x < 50, 50 < x < 60, 60 < x < 70, 70 < x < 80, 80 < x < 90, 90 < x < 100$ ”.

Estimation of similarity values and ranking of the samples

The values of the MF of each sample, i.e., B_1, B_2, B_3 , and B_4 were compared with the values of the MF of the standard scale, i.e., F_1, F_2, F_3, F_4, F_5 , and F_6 given in equation 7. Each of the terms $B_1, B_2, B_3, B_4, F_1, F_2, F_3, F_4, F_5$, and F_6 represent a row matrix having 10 elements.

The similarity values of the samples B were calculated using equation 9 [44]. Afterward, the similarity values of four samples were compared, and then the sample with the higher similarity value was considered extremely important (EI):

$$S_m(F_j, B_x) = \frac{F_j \times B_x^T}{\text{Max}(F_j \times F_j^T \text{ and } B_x \times B_x^T)}$$

Where S_m represents the similarity value of the sample B_x ; F_j represents F_1, F_2, F_3, F_4, F_5 ; B_x is the MF value for sample x on the standard fuzzy scale; B_x^T and F_j^T are transposes of the matrix B_x and F_j . For example, for functional extruded samples (S_1): $S_m(F_1, B_1), S_m(F_2, B_1), S_m(F_3, B_1), S_m(F_4, B_1), S_m(F_5, B_1), S_m(F_6, B_1)$ were calculated from equation 9 using the matrix multiplication rule. Similarly, similarity values of other commercial samples S_2, S_3 , and S_4 were calculated. Similarity values under the 6-scale factors of standard fuzzy sensory scales are compared to find out the highest similarity value. The same procedure was followed to rank the quality attributes (color, flavor, taste, and mouthfeel) of the four samples in general.

Estimation of similarity values and ranking of quality attributes for each sample

A similar procedure was followed to find out the ranks of four quality attributes for each of the samples S_1, S_2, S_3 , and S_4 . The only difference is that Q_{sum} is obtained by calculating the average of the first digit of the triplets of QC, QA, QT , and QM .

Results

PR

Phytosterols are sterols derived from plants, similar to cholesterol in terms of structure. Their degradation mainly depends on factors including temperature, moisture, light, processing conditions, and storage.

Various studies have shown that high-temperature processing techniques like heating, baking, and frying result in phytosterols' degradation and form various free radicals. In this study, the PR of the extrudates ranged from 0.4832 g/100 g to 0.9532 g/100 g. The highest PR was observed at an extrusion temperature of 130°C, whereas the lowest was observed at a temperature of 150°C. This might be because phytosterols were unstable/degraded and oxidized at elevated temperature, generating free radicals [45]. In another study, vitamin B₁₂ in extrudates degraded at higher temperatures and lost completely at 190°C during the extrusion process [46]. Various food products fortified with phytosterol including cheese spread from goat milk, functional fermented maize yogurt, and fermented cereal products showed degradation of phytosterols at elevated extrusion temperature [16, 47, 48].

Overall SS of the samples

The sum of the SS given by panelists/judges and the triplets associated to the quality attributes of different samples are given in Table 3 and sum of the SS for the quality attributes for the samples in general is given in Table 4. The relative attributes of the samples were measured using five-point sensory scale factors. Sensory scale factors were used to calculate general quality attributes like NI, SI, important (I), highly important (HI), and EI. Next, the quality attributes and relative weights were determined using equation 3. Finally, the overall scores of the samples were estimated by multiplying the triplets of quality attributes of the samples and relative weights of quality attributes of the samples using the multiplication rule as given in equation 6. The overall SS of individual samples calculated in the form of triplets are given below:

$$SO_1 = (51.1759 \ 40.9537 \ 34.2500)$$

$$SO_2 = (73.8889 \ 49.5833 \ 38.1296)$$

$$SO_3 = (72.9722 \ 48.9722 \ 33.2407)$$

$$SO_4 = (74.5556 \ 50.0000 \ 33.1481)$$

Where SO_1 , SO_2 , SO_3 , and SO_4 are the overall SS of the samples S_1 , S_2 , S_3 , and S_4 , respectively.

Overall MF of the samples on the fuzzy logic scale

The values and symbols of the six-point scale MF are shown in equation 7.

The value of the MF of the overall SS of each sample on the standard fuzzy scale at $x = 0$ to 100 were calculated using equation 8, and the values are as shown below:

$$B_1 = (0, 0.2388, 0.4829, 0.7271, 0.9713, 1, 0.7424, 0.4504, 0.1584, 0)$$

$$B_2 = (0, 0, 0.1148, 0.3165, 0.5182, 0.7199, 0.9216, 1, 0.8397, 0.5775)$$

$$B_3 = (0, 0, 0.1225, 0.3267, 0.5309, 0.7351, 0.9393, 1, 0.7886, 0.4877)$$

$$B_4 = (0, 0, 0.1089, 0.3089, 0.5089, 0.7089, 0.9089, 1, 0.8358, 0.5341)$$

Similarity values and the ranking of the samples

The similarity values (S_m) and the ranking of the food products indicate the comparative importance of all the quality characteristics that aid in acceptance/rejection. Similarity values of the samples were calculated by combining the MF values of the standard fuzzy logic scale (F 's) and the overall MF values of the SS (B 's) using equation 9 and the calculated values are shown in Table 5. For example, the similarity values of S_1 were calculated as 0, 0.0337, 0.3064, 0.6890, 0.6927, 0.2767, and 0.0224, which were not satisfactory, fair, satisfactory, good, very good, and excellent respectively. Thus, the overall SS of S_1 was good due to its high S_m value. Similarly, S_2 , S_3 , and S_4 were very good with S_m values of 0.6835, 0.6788, and 0.6952, respectively as shown in Table 5.

Table 5. Similarity values of the samples under various scale factors

Scale factor	Similarity values (S_m)			
	S_1	S_2	S_3	S_4
Not satisfactory, F_1	0.0337	0	0	0
Fair, F_2	0.3064	0.0721	0.0776	0.0716
Satisfactory, F_3	0.6890	0.3305	0.3490	0.3335
Good, F_4	0.6927	0.6337	0.6619	0.6449
Very good, F_5	0.2767	0.6835	0.6788	0.6952
Excellent, F_6	0.0224	0.2633	0.2393	0.2588

Similarity values and ranking of quality attributes of samples in general

Similar procedure was used for ranking of quality attributes (color, flavor, taste, and mouthfeel) for the evaluation of the samples in general. Triplets of overall SS for quality attributes were calculated in equation 3. Using the method of computation mentioned earlier and denoting BC , BA , BT , and BM as overall SS on the standard fuzzy scale, for color, flavor, taste, and mouthfeel, respectively we get:

$$BC = (0, 0, 0, 0, 0.1333, 0.5333, 0.9333, 1, 0.5455, 0)$$

$$BA = (0, 0, 0, 0.2, 0.6, 1, 1, 0.5385, 0.0769, 0)$$

$$BT = (0, 0, 0, 0, 0, 0.1333, 0.5333, 0.9333, 1, 0.5)$$

$$BM = (0, 0, 0, 0, 0, 0, 0.3333, 0.7333, 1, 0.7143)$$

Similarity values, S_m of all the four quality attributes (color, flavor, taste, and mouthfeel) were calculated and are presented in Table 6.

Table 6. Similarity values of quality attributes of samples in general

Scale factor	Color	Flavor	Taste	Mouthfeel
Not at all necessary, F_1	0	0	0	0
Somewhat necessary, F_2	0	0.0371	0	0
Necessary, F_3	0.1600	0.4822	0.0267	0
I, F_4	0.8133	0.9530	0.4533	0.2800
HI, F_5	0.8048	0.4137	0.9800	0.9029
EI, F_6	0.1104	0.0143	0.4127	0.5624

Similarity values and the ranking for each quality attribute of the samples

Similarity values of the four quality attributes of each sample against six-point sensory scales, i.e., not satisfactory, fair, satisfactory, good, very good, and excellent were calculated using equations 7, 8, and 9. Using the method of computation mentioned earlier and denoting BC_1 , BA_1 , BT_1 , and BM_1 as overall SS on standard fuzzy scale, for color, flavor, taste, and mouthfeel, respectively, for S_1 we get:

$$BC_1 = (0.1708, 0.4646, 0.7584, 1, 0.9472, 0.6502, 0.3531, 0.0561, 0, 0)$$

$$BA_1 = (0.0379, 0.3003, 0.5627, 0.8251, 1, 0.8985, 0.5941, 0.2897, 0, 0)$$

$$BT_1 = (0, 0.1842, 0.4092, 0.6342, 0.8592, 1, 0.8986, 0.6275, 0.3564, 0.0853)$$

$$BM_1 = (0, 0.0776, 0.2894, 0.5012, 0.7129, 0.9247, 1, 0.8082, 0.5105, 0.2128)$$

Similarly, overall SS for all the quality attributes of S_2 , S_3 , and S_4 were calculated and are given below.

For S_2 :

$$BC_2 = (0, 0, 0.1030, 0.3030, 0.5030, 0.7030, 0.9030, 1, 0.8654, 0.6041)$$

$$BA_2 = (0, 0.1692, 0.4000, 0.6308, 0.8615, 1, 0.8896, 0.6135, 0.3374, 0.0613)$$

$$BT_2 = (0, 0, 0.0601, 0.2558, 0.4514, 0.6471, 0.8428, 1, 0.9525, 0.7106)$$

Table 3. Sum of SS for the quality attributes of the samples and its associated triplets

Sensory attributes	Samples	Not satisfactory	Fair	Medium	Good	Excellent	Triplets associated with the sensory scale
Color	S_1	2	3	9	1	0	(40.00 21.67 25.00)
	S_2	0	0	1	11	3	(78.33 25.00 20.00)
	S_3	0	0	3	3	9	(85.00 25.00 10.00)
	S_4	0	1	1	3	10	(86.67 25.00 8.33)
Flavor	S_1	1	4	2	5	3	(58.33 23.33 20.00)
	S_2	0	1	4	7	3	(70.00 25.00 20.00)
	S_3	1	2	1	8	3	(66.67 23.33 20.00)
	S_4	0	0	4	7	4	(75.00 25.00 18.33)
Taste	S_1	0	4	7	3	1	(51.67 25.00 23.33)
	S_2	0	0	3	11	1	(71.67 25.00 23.33)
	S_3	0	3	3	5	4	(66.67 25.00 18.33)
	S_4	0	2	5	4	4	(66.67 25.00 18.33)
Mouthfeel	S_1	0	4	6	3	2	(55.00 25.00 21.67)
	S_2	0	0	2	11	2	(75.00 25.00 21.67)
	S_3	0	1	3	7	4	(73.33 25.00 18.33)
	S_4	0	1	4	6	4	(71.67 25.00 18.33)

Table 4. Sum of SS for the quality attributes of the samples in general and triplets associated with the SS and relative weights of the quality attributes of the samples in general

Quality attribute	NI	SI	I	HI	EI	Triplets	RW
						SS	
Color	0	0	6	5	4	(71.67 25.00 18.33)	(0.24 0.08 0.06)
Flavor	0	3	5	5	2	(60.00 25.00 21.67)	(0.20 0.08 0.07)
Taste	0	0	1	9	5	(81.67 25.00 16.68)	(0.27 0.08 0.06)
Mouthfeel	0	0	1	6	8	(86.67 25.00 11.68)	(0.29 0.08 0.039)

RW: relative weightage

$$BM_2 = (0, 0, 0, 0.1340, 0.3196, 0.5052, 0.6907, 0.8763, 1, 0.9092)$$

For S_3 :

$$BC_3 = (0, 0, 0.0191, 0.2106, 0.4021, 0.5936, 0.7851, 0.9766, 1, 0.7106)$$

$$BA_3 = (0, 0.1848, 0.4293, 0.6739, 0.9185, 1, 0.8109, 0.5273, 0.2437, 0)$$

$$BT_3 = (0, 0, 0.1386, 0.3408, 0.5431, 0.7453, 0.9476, 1, 0.7870, 0.4995)$$

$$BM_3 = (0, 0, 0, 0.1611, 0.3486, 0.5361, 0.7236, 0.9111, 1, 0.8386)$$

For S_4 :

$$BC_4 = (0, 0, 0, 0.1888, 0.3782, 0.5677, 0.7572, 0.9467, 1, 0.7535)$$

$$BA_4 = (0, 0.1111, 0.3333, 0.5556, 0.7778, 1.0000, 1, 0.7248, 0.4495, 0.1743)$$

$$BT_4 = (0, 0, 0.1386, 0.3408, 0.5431, 0.7453, 0.9476, 1, 0.7870, 0.4995)$$

$$BM_4 = (0, 0, 0, 0.1888, 0.3782, 0.5677, 0.7572, 0.9467, 1, 0.7778)$$

The similarity values of each quality attribute of the samples S_1 , S_2 , S_3 , and S_4 are shown in [Table 7](#).

Table 7. Similarity values for the quality attributes of all the samples S_1 , S_2 , S_3 , and S_4

Sensory scale	Color	Flavor	Taste	Mouthfeel
Functional extrudate (S_1)				
Not satisfactory, F_1	0.1234	0.0564	0.0250	0.0106
Fair, F_2	0.5534	0.3883	0.2476	0.1686
Satisfactory, F_3	0.8113	0.7667	0.5977	0.4971
Good, F_4	0.4605	0.6413	0.7185	0.7330

Table 7. Similarity values for the quality attributes of all the samples S_1 , S_2 , S_3 , and S_4 (continued)

Sensory scale	Color	Flavor	Taste	Mouthfeel
Very good, F_5	0.0712	0.1760	0.4014	0.5255
Excellent, F_6	0	0	0.0717	0.1278
Commercial sample (S_2)				
Not satisfactory, F_1	0	0.0234	0	0
Fair, F_2	0.0673	0.2448	0.0493	0.0194
Satisfactory, F_3	0.3199	0.6066	0.2782	0.2049
Good, F_4	0.6239	0.7269	0.5809	0.5204
Very good, F_5	0.6931	0.3947	0.7156	0.7764
Excellent, F_6	0.2744	0.0637	0.3112	0.4088
Commercial sample (S_3)				
Not satisfactory, F_1	0	0.0263	0	0
Fair, F_2	0.0343	0.2709	0.0823	0.0231
Satisfactory, F_3	0.2529	0.6570	0.3533	0.2227
Good, F_4	0.5691	0.7216	0.6567	0.5411
Very good, F_5	0.7497	0.3351	0.6690	0.7710
Excellent, F_6	0.3331	0.0347	0.2380	0.3833
Commercial sample (S_4)				
Not satisfactory, F_1	0	0.0146	0	0
Fair, F_2	0.0267	0.1903	0.0823	0.0264
Satisfactory, F_3	0.2405	0.5270	0.3533	0.2380
Good, F_4	0.5617	0.7250	0.6567	0.5558
Very good, F_5	0.7637	0.4642	0.6690	0.7591
Excellent, F_6	0.3543	0.1052	0.2380	0.3574

Discussion

From Table 5, it was observed that S_4 had the maximum similarity value of 0.6952 (very good category) among all the samples. S_2 and S_3 were also found in the “very good” category with similarity values of 0.6835 and 0.6788, and S_1 was in the “good” category with a value of 0.6927. The ranking order for the extrudate samples is as follows: S_4 (very good) > S_2 (very good) > S_3 (very good) > S_1 (good), i.e., commercial sample (S_4) was the most accepted extruded sample by all panelists/judges followed by S_2 , S_3 , and S_1 .

From Table 6, it was observed that compared to taste and mouthfeel, color and flavor have less impact on assessing the quality attributes which fell under the I category. Taste and mouthfeel were under the HI category. These results led to the following general rankings of quality attributes for the samples: taste (HI) > mouthfeel (HI) > flavor (I) > color (I).

For functional extrudate sample (S_1), S_m values for mouthfeel and taste were found to be 0.7330 and 0.7185 respectively under “good” category while for color and flavor these values were 0.8113 and 0.7667, respectively, and were under “satisfactory” category. Therefore, taste is the strongest quality of functional extrudate sample (S_1) while its flavor is the weakest. Hence, improvement in the flavor of the functional extrudates can help to increase its marketability. For the other three commercial samples (S_2 , S_3 , and S_4), color, taste, and mouthfeel fall under “very good” category while flavor falls under “good” category.

Conclusions

The sensory characteristics of functional extrudates were assessed based on their quality attributes, using the fuzzy logic technique. Before the sensory analysis was conducted, functional extrudates were developed by twin screw extrusion process of phytosterol enriched corn and pea protein. Functional extrudates (S_1) were compared with commercial samples (S_2 , S_3 , and S_4). Among these, S_4 achieved the top rank, with a sensory scale labeled as “very good” and a similarity value of 0.6952. It was followed by S_2 , then S_3 , and finally the functional extrudates (S_1). The overall SS for the samples, as obtained from the similarity values are:

S_1 : mouthfeel (good) > taste (good) > color (satisfactory) > flavor (satisfactory)
 S_2 : mouthfeel (very good) > taste (very good) > color (very good) > flavor (good)
 S_3 : mouthfeel (very good) > color (very good) > taste (very good) > flavor (good)
 S_4 : color (very good) > mouthfeel (very good) > taste (very good) > aroma (good)

Abbreviations

El: extremely important

HI: highly important

I: important

MF: membership function

NI: not at all important

PPI: pea protein isolates

PR: phytosterol retention

SI: somewhat important

SS: sensory scores

Declarations

Acknowledgments

The authors are grateful to the Centralized Research Facility (CRF) of NIT Rourkela for allowing them to use analytical instruments.

Author contributions

MP: Writing—original draft. PS: Conceptualization, Formal analysis, Validation, Writing—review & editing. DTR: Writing—review & editing. SKS: Methodology, Formal analysis, Validation, Conceptualization, Funding acquisition, Supervision, Project administration, Writing—review & editing.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Ethical approval

Not applicable.

Consent to participate

Not applicable.

Consent to publication

Not applicable.

Availability of data and materials

Datasets are available from the corresponding author upon reasonable request.

Funding

This work is funded by the Science and Engineering Research Board [SRG/2019/000998], New Delhi, India. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Singh SK, Muthukumarappan K. Effect of different extrusion processing parameters on physical properties of soy white flakes and high protein distillers dried grains-based extruded aquafeeds. *J Food Res.* 2014;3:107–23.
2. Singh SK, Muthukumarappan K. Single screw extrusion processing of soy white flakes based catla feed. *J Food Res.* 2015;4:1–9.
3. Singh SK, Muthukumarappan K. Effect of feed moisture, extrusion temperature and screw speed on properties of soy white flakes based aquafeed: a response surface analysis. *J Sci Food Agric.* 2016;96:2220–9.
4. Singh SK. Understanding the effect of extrusion processing parameters on physical, nutritional and rheological properties of soy white flakes based aquafeed in a single screw extruder [dissertation]. Ann Arbor: South Dakota State University; 2016. p. 252.
5. Singha P. Understanding the impact of extrusion processing on rheological, textural and functional properties of high-protein, high-fiber extrudates [dissertation]. Ann Arbor: South Dakota State University; 2017. p. 313.
6. Singha P, Singh SK, Muthukumarappan K. Textural and structural characterization of extrudates from apple pomace, defatted soy flour and corn grits. *J Food Process Eng.* 2019;42:e13046.
7. Singha P, Singh SK, Muthukumarappan K, Krishnan P. Physicochemical and nutritional properties of extrudates from food grade distiller's dried grains, garbanzo flour, and corn grits. *Food Sci Nutr.* 2018;6:1914–26.
8. Yoon AK, Singha P, Rizvi SSH. Steam vs. SC-CO₂-based extrusion: comparison of physical properties of milk protein concentrate extrudates. *J Food Eng.* 2021;292:110244.
9. Rostamabadi H, Bajer D, Demirkesen I, Kumar Y, Su C, Wang Y, et al. Starch modification through its combination with other molecules: gums, mucilages, polyphenols and salts. *Carbohydr Polym.* 2023;314:120905.
10. Ziena HM, Ziena AHM. Nutritious novel snacks from some of cereals, legumes and skimmed milk powder. *Appl Food Res.* 2022;2:100092.
11. Dash DR, Singh SK, Singha P. Recent advances on the impact of novel non-thermal technologies on structure and functionality of plant proteins: a comprehensive review. *Crit Rev Food Sci Nutr.* 2022;1–16.
12. Asaithambi N, Singha P, Singh SK. Recent application of protein hydrolysates in food texture modification. *Crit Rev Food Sci Nutr.* 2023;63:10412–43.
13. Barbhuiya RI, Singh SK, Singha P. Mangosteen wastes. In: Muzaffar K, Ahmad Sofi S, Mir SA, editors. *Handbook of fruit wastes and by-products. Chemistry, processing technology, and utilization.* Boca Raton: CRC Press; 2022. pp. 113–24.
14. Barbhuiya RI, Singha P, Singh SK. Pomelo Wastes. In: Muzaffar K, Ahmad Sofi S, Mir SA, editors. *Handbook of fruit wastes and by-products. Chemistry, processing technology, and utilization.* Boca Raton: CRC Press; 2022. pp. 19–38.
15. Shekhar S, Prakash P, Singha P, Prasad K, Singh SK. Modeling and optimization of ultrasound-assisted extraction of bioactive compounds from *Allium sativum* leaves using response surface methodology and artificial neural network coupled with genetic algorithm. *Foods.* 2023;12:1925.
16. Amaral JBS, Grisi CVB, Vieira EA, Ferreira PS, Rodrigues CG, Diniz NCM, et al. Light cream cheese spread of goat milk enriched with phytosterols: physicochemical, rheological, and microbiological characterization. *LWT.* 2022;157:113103.

17. Yuan Y, Liu C, Zhao G, Gong X, Dang K, Yang Q, et al. Transcriptome analysis reveals the mechanism associated with dynamic changes in fatty acid and phytosterol content in foxtail millet (*Setaria italica*) during seed development. *Food Res Int.* 2021;145:110429.
18. Feng S, Wang L, Shao P, Lu B, Chen Y, Sun P. Simultaneous analysis of free phytosterols and phytosterol glycosides in rice bran by SPE/GC–MS. *Food Chem.* 2022;387:132742.
19. Guo SJ, Ma CG, Hu YY, Bai G, Song ZJ, Cao XQ. Solid lipid nanoparticles for phytosterols delivery: the acyl chain number of the glyceride matrix affects the arrangement, stability, and release. *Food Chem.* 2022;394:133412.
20. Jakubczyk E, Linde M, Gondek E, Kamińska-Dwórznińska A, Samborska K, Antoniuk A. The effect of phytosterols addition on the textural properties of extruded crisp bread. *J Food Eng.* 2015;167:156–61.
21. Gao K, Rao J, Chen B. Unraveling the mechanism by which high intensity ultrasound improves the solubility of commercial pea protein isolates. *Food Hydrocolloids.* 2022;131:107823.
22. Li N, Yang M, Guo Y, Tong LT, Wang Y, Zhang S, et al. Physicochemical properties of different pea proteins in relation to their gelation ability to form lactic acid bacteria induced yogurt gel. *LWT.* 2022;161:113381.
23. Beck SM, Knoerzer K, Foerster M, Mayo S, Philipp C, Arcot J. Low moisture extrusion of pea protein and pea fibre fortified rice starch blends. *J Food Eng.* 2018;231:61–71.
24. Philipp C, Emin MA, Buckow R, Silcock P, Oey I. Pea protein-fortified extruded snacks: linking melt viscosity and glass transition temperature with expansion behaviour. *J Food Eng.* 2018;217:93–100.
25. Philipp C, Oey I, Silcock P, Beck SM, Buckow R. Impact of protein content on physical and microstructural properties of extruded rice starch-pea protein snacks. *J Food Eng.* 2017;212:165–73.
26. Singh SK, Muthukumarappan K. Rheological characterization and CFD simulation of soy white flakes based dough in a single screw extruder. *J Food Process Eng.* 2017;40:e12368.
27. Singha P, Muthukumarappan K, Krishnan P. Influence of processing conditions on apparent viscosity and system parameters during extrusion of distiller's dried grains-based snacks. *Food Sci Nutr.* 2018;6:101–10.
28. Singha P, Muthukumarappan K. Single screw extrusion of apple pomace-enriched blends: Extrudate characteristics and determination of optimum processing conditions. *Food Sci Technol Int.* 2018;24:447–62.
29. Singha P, Muthukumarappan K. Effects of processing conditions on the system parameters during single screw extrusion of blend containing apple pomace. *J Food Process Eng.* 2017;40:e12513.
30. Singh SK, Singha P, Muthukumarappan K. Modeling and optimizing the effect of extrusion processing parameters on nutritional properties of soy white flakes-based extrudates using response surface methodology. *Anim Feed Sci Technol.* 2019;254:114197.
31. Singha P, Muthukumarappan K. Quality changes and freezing time prediction during freezing and thawing of ginger. *Food Sci Nutr.* 2016;4:521–33.
32. Singh SK, Muthukumarappan K. A viscosity model for soy white flakes-based aquafeed dough in a single screw extruder. *J Food Process Eng.* 2017;40:e12357.
33. Basak S. The use of fuzzy logic to determine the concentration of betel leaf essential oil and its potency as a juice preservative. *Food Chem.* 2018;240:1113–20.
34. Mukhopadhyay S, Majumdar GC, Goswami TK, Mishra HN. Fuzzy logic (similarity analysis) approach for sensory evaluation of *chhana podo*. *LWT.* 2013;53:204–10.
35. Sugumar JK, Guha P. Comparative study on the hedonic and fuzzy logic based sensory analysis of formulated soup mix. *Future Foods.* 2022;5:100115.
36. Das S, Pegu K, Arya SS. Functional sourdough millet bread rich in dietary fibre -an optimization study using fuzzy logic analysis. *Bioact Carbohydr Diet Fibre.* 2021;26:100279.

37. Kumar R, Ghosh P, Srinivasa Rao P, Rana SS, Vashishth R, Vivek K. Sensory evaluation of microwave assisted ultrasound treated soymilk beverage using fuzzy logic. *J Saudi Soc Agric Sci.* 2021;20:257–64.
38. Jaya S, Das H. Sensory evaluation of mango drinks using fuzzy logic. *J Sens Stud.* 2003;18:163–76.
39. Kaushik N, Gondi AR, Rana R, Srinivasa Rao P. Application of fuzzy logic technique for sensory evaluation of high pressure processed mango pulp and litchi juice and its comparison to thermal treatment. *Innovative Food Sci Emerging Technol.* 2015;32:70–8.
40. Tahsiri Z, Niakousari M, Khoshnoudi-Nia S, Hosseini SMH. Sensory evaluation of selected formulated milk barberry drinks using the fuzzy approach. *Food Sci Nutr.* 2017;5:739–49.
41. Pavani M, Singha P, Singh SK. Development of phytosterol enriched functional edible oils: study of physical, chemical, thermal and structural Properties. *J Sci Ind Res.* 2022;81:549–60.
42. Pavani M, Singha P, Rajamanickam DT, Singh SK. Impact of extrusion processing on bioactive compound enriched plant-based extrudates: a comprehensive study and optimization using RSM and ANN-GA. *Future Foods.* 2023;100286.
43. Araújo LBDC, Silva SL, Galvão MAM, Ferreira MRA, Araújo EL, Randau KP, et al. Total phytosterol content in drug materials and extracts from roots of *Acanthospermum hispidum* by UV-VIS spectrophotometry. *Rev Bras Farmacogn.* 2013;23:736–42.
44. Das H. Sensory evaluation using fuzzy logic. In: *Food processing operations analysis.* New Delhi: Asian Books Pvt. Ltd.; 2005. pp. 383–402.
45. Pavani M, Singha P, Dash DR, Asaithambi N, Singh SK. Novel encapsulation approaches for phytosterols and their importance in food products: a review. *J Food Process Eng.* 2022;45:e14041.
46. Bajaj SR, Singhal RS. Effect of extrusion processing and hydrocolloids on the stability of added vitamin B₁₂ and physico-functional properties of the fortified puffed extrudates. *LWT.* 2019;101:32–9.
47. Gies M, Descalzo AM, Servent A, Dhuique-Mayer C. Incorporation and stability of carotenoids in a functional fermented maize yogurt-like product containing phytosterols. *LWT.* 2019;111:105–10.
48. Lingua MS, Gies M, Descalzo AM, Servent A, Páez RB, Baroni MV, et al. Impact of storage on the functional characteristics of a fermented cereal product with probiotic potential, containing fruits and phytosterols. *Food Chem.* 2022;370:130993.